

lipid synthesis, nevertheless, it suggests that in the early stages of maturation the cell is accumulating protein more rapidly than lipid. It is possible, therefore, that at the onset of differentiation the cell density does not decrease but actually increases. Our attempts to separate undifferentiated from early differentiating cells have been unsuccessful presumably because, for the above reasons, the density difference is too small.

The second observation refers to the changes in lipid composition of the tumor cells as they decrease in buoyant density. Similar changes have been observed in preliminary experiments with normal cells so this is not a feature unique to the tumor cell. Studies of the skin surface lipids of man by Downing and Strauss [25,26] have indicated that changes may occur in the lipid composition of the sebaceous cell as it matures and this is also implied in the studies on isolated human sebaceous glands by Summerly [27], but direct evidence has been lacking. These observations are of interest since they suggest that sequential gene activation may be occurring during the maturation of the sebaceous cell and that the term cytodifferentiation may, justifiably, be used to describe this process.

Even though less extensive studies have been made with normal gland cells, nevertheless the evidence so far available shows distinct differences from the tumor cells. Hence, while the tumor enables large numbers of cells to be easily collected for use in biochemical investigations, the data from such studies will need to be interpreted with caution until it can be confirmed on normal cells. Further studies with normal cells will define more clearly the uses and limitations of the tumor cells in the study of sebaceous cell differentiation. The cells obtained from both the tumor and the normal gland by these procedures are viable cells which are suitable for chemical and biochemical assays of relatively short duration. They are not intended for long-term incubation studies for which the cloned cell lines previously described by us [28] would be more suitable. Their advantage over the cultured cells is that they represent cells as they exist in the parent tissue and have not been exposed to an artificial environment as occurs with cells maintained in culture.

REFERENCES

1. Bizzozero G, Vassale G: Ueber die Erzeugen und die physiologische Regeneration der Drüsenzellen bei Säugethieren. *Virchows Arch A* 110:155-214, 1887
2. Ellis RA, Hendrickson RG: The ultrastructure of the sebaceous gland of man. *Adv Biol Skin* 4:94-109, 1963
3. Bell M: A comparative study of the ultrastructure of the sebaceous glands of man and other primates. *J Invest Dermatol* 62:132-143, 1974
4. Gill GW: Exfoliative cytology. Millipore Application Report AR-24, 1970
5. Lowry OH, Rosebrough NJ, Farr AL, Randall RJ: Protein measurement with the Folin phenol reagent. *J Biol Chem* 193:265-275, 1951
6. Vorbeck ML, Martinetti GV: Separation of glycosyl diglycerides from phosphatides using silicic acid column chromatography. *J Lipid Res* 6:3-6, 1965
7. Bloor WR: A colorimetric procedure for the determination of small amounts of fatty acid. *J Biol Chem* 170:671-674, 1947
8. Downing DT: Photodensitometry in the thin-layer chromatographic analysis of neutral lipids. *J Chromatog* 38:91-99, 1968
9. Halprin KM, Ohkawara A: Glucose and glycogen metabolism in the human epidermis. *J Invest Dermatol* 46:43-50, 1966
10. Im MJC, Hoopes JE: Enzymes of carbohydrate metabolism in normal human sebaceous glands. *J Invest Dermatol* 62:153-160, 1974
11. Srere PA: Citrate-cleavage enzyme. *Methods Enzymol* 5:641-642, 1962
12. Numa S: Acetyl-CoA carboxylase from chicken and rat liver. *Methods Enzymol* 14:9-10, 1969
13. Everhart LP, Hauschka PV, Prescott DM: Measurement of growth and rates of incorporation of radioactive precursors into macromolecules of cultured cells. *Methods Cell Biol* 7:329-346, 1973
14. Bush ET: General applicability of the channels ratio method for measuring liquid scintillation efficiencies. *Anal Chem* 35:1024-1029, 1963
15. Wheatley VR, Hodgins LT, Coon WM, Kumarasiri M, Berenzweig H, Feinstein JM: Cutaneous lipogenesis. Precursors utilized by guinea pig ear skin for lipid synthesis. *J Lipid Res* 12:347-360, 1971
16. Strong LC: The transplantation of an adenocarcinoma of the preputial gland in mice of the A strain. *Cancer Res* 2:332-334, 1942
17. Brown JC, Williams JA: A histochemical study of the preputial glands of the male laboratory rat and mouse. *Acta Anat* 81:270-285, 1972
18. Prutkin L, Rossman TG, Wheatley VR: Ultrastructural changes and viral morphogenesis during the growth of the mouse preputial gland tumor ESR 586. *Cancer Res* 37:551-556, 1977
19. Fidler IJ: Immune stimulation-inhibition of experimental cancer metastasis. *Cancer Res* 34:491-498, 1974
20. Bellemann P, Gebhardt R, Mecke D: An improved method for the isolation of hepatocytes from liver slices. *Anal Biochem* 81:408-415, 1977
21. deOca HM: Trypsin B. High yield method for kidney tissue, *Tissue Culture Methods and Applications*. Edited by PF Kruse, MK Patterson, New York, Academic Press, pp 8-12, 1973
22. Shall S: Cells at defined stages of the cell cycle, *Tissue Culture Methods and Applications*. Edited by PF Kruse, MK Patterson, New York, Academic Press, pp 195-204, 1973
23. Tosti A: A comparison of the histodynamics of sebaceous glands and epidermis in man: A microanatomic and morphometric study. *J Invest Dermatol* 62:147-152, 1974
24. Beaver DL: A re-evaluation of the rat preputial gland as a "dicrine" organ from the standpoint of morphology, histochemistry and physiology. *J Exp Zool* 143:153-173, 1960
25. Downing DT, Strauss JS: Synthesis and composition of surface lipids of human skin. *J Invest Dermatol* 62:228-244, 1974
26. Downing DT, Strauss JS, Norton LA, Pochi PE, Stewart ME: The time course of lipid formation in human sebaceous glands. *J Invest Dermatol* 69:407-412, 1977
27. Summerly R: Skin components: ^{14}C acetate incorporation in vitro, with particular reference to lipid synthesis in isolated (dissected) human sebaceous glands. *Trans St Johns Hospital Derm Soc* 60: 1-23, 1974
28. Rossman TG, Prutkin L, Wheatley VR: Culture of cloned cells from the mouse preputial gland tumor ESR 586. *Cancer Res* 37:557-560, 1977

Announcement

The Fourth Conference on Cutaneous Toxicity sponsored by the American Medical Association and the Society of Toxicology will be held at the Mayflower Hotel, Washington, D.C., May 9-11th, 1979. This continuing medical educational activity is acceptable for 12 credit hours in Category No. 2 for the Physician's Recognition Award of the American Medical Association. Registration fee \$150 (\$120 for AMA and SOT members; \$85 for residents and retired physicians). For further information contact Dr. Joseph B. Jerome, American Medical Association, 535 North Dearborn Street, Chicago, Illinois 60601.